26 July 1999

UMN-TH-1803/99
UMN-TH-1810/99

arXiv:gr-qc/9907091v2 9 Aug 1999

Monopole and dyon solutions in the Einstein Yang-Mills theory in asymptotically anti-de Sitter space*

Yutaka Hosotani and Jefferson Bjoraker

School of Physics and Astronomy, University of Minnesota
Minneapolis, MN 55455, U.S.A.

Abstract

Regular monopole and dyon solutions to the $SU(2)$ Einstein Yang-Mills equations in asymptotically anti-de Sitter space are discussed. A class of monopole solutions are shown to be stable against spherically symmetric linear perturbations.

1 Introduction

Static solutions to the Einstein Yang-Mills (EYM) equations differ considerably depending on the value of the cosmological constant. The solutions can be separated into two families; $\Lambda \geq$

*Proceedings from the talk given at "The eighth Canadian Conference on General Relativity and Astrophysics", McGill University, Montreal, Quebec, Canada, June 10-12 1999.

0, and $\Lambda < 0$. The solutions where $\Lambda = 0$ were discovered by Bartnik and McKinnon (BK) [1] and their asymptotically de Sitter (dS) analogs ($\Lambda > 0$) were discovered independently by Volkov et. al. and Torii et. al. [2]. The BK solutions and the cosmological extensions to them all share similar behavior and have been studied in detail (see Ref. [3] for a review). Recently, asymptotically anti-de Sitter ($\Lambda < 0$) black hole solutions [4] and soliton solutions [5] were found which are strikingly different than the BK type solutions. In particular, there exist asymptotically anti-de Sitter (AdS) solitons with no nodes in the field strength which are stable. Furthermore, in asymptotically AdS space dyon solutions are allowed.

2 General formalism

Given the spherically symmetric metric in the Schwarzschild gauge

$$ds^2 = -\frac{H dt^2}{p^2} + \frac{dr^2}{H} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \tag{1}$$

where H, p are functions of t and r, the coupled static EYM equations of motion are

$$\left(\frac{H}{p}w'\right)' = -\frac{p}{H}u^2 w - \frac{w}{p}\frac{(1-w^2)}{r^2} \tag{2}$$

$$\left(r^2 p u'\right)' = \frac{2p}{H}w^2 u \tag{3}$$

$$p' = -\frac{2v}{r}p\left[(w')^2 + \frac{u^2 w^2 p^2}{H^2}\right] \tag{4}$$

$$m' = v\left[\frac{(w^2-1)^2}{2r^2} + \frac{1}{2}r^2 p^2 (u')^2 + H(w')^2 + \frac{u^2 w^2 p^2}{H}\right], \tag{5}$$

where $w(r)$ and $u(r)$ are the magnetic and electric components of the $SU(2)$ Yang-Mills fields [6], $H(r) = 1 - 2m(r)/r - \Lambda r^2/3$, $v = G/4\pi e^2$ and Λ is the cosmological constant. We require that solutions to Eq.'s (2) to (5) are regular everywhere and have finite ADM mass $M = m(\infty)$. The electric and magnetic charges of solutions are given by

$$\begin{pmatrix} Q_E \\ Q_M \end{pmatrix} = \frac{e}{4\pi}\int dS_k \sqrt{-g}\begin{pmatrix} F^{k0} \\ \tilde{F}^{k0} \end{pmatrix} = \begin{pmatrix} u_1 p_0 \\ 1 - w_0^2 \end{pmatrix}\frac{\tau_3}{2} \tag{6}$$

where $w = w_0 + w_1/r + \mathrm{O}(r^{-2})$ etc.

For solutions in asymptotically flat or dS space, $w(r)$ has at least one node [2]. The situation is quite different in asymptotically AdS space ($\Lambda < 0$). $H(r)$ is positive everywhere and there are solutions where w has no node.

3 Monopole solutions

The solutions to Eq.'s (2) to (5), for $\Lambda < 0$, were evaluated numerically [5] using the shooting method. In the shooting method one solves Eqs. (2) to (5) at $r = 0$ in terms of two parameters, a and b, and 'shoots' for solutions with the desired asymptotic behavior. a and b are adjustable parameters which together specify the boundary conditions at the origin for w, u, H, and p: $u(r) = ar + \cdots$ and $w(r) = 1 - br^2 + \cdots$ near $r = 0$.

Purely magnetic solutions (monopoles) are found by setting $a = 0$, corresponding to $u(r) = 0$. A continuum of monopole solutions were obtained by varying the parameter b. The solutions are similar to the black hole solutions found by Ref. [4], but are also regular for all r. The number of times w crosses the axis depends on the value of the adjustable shooting parameter b.

As shown in fig. 1, the behavior of m and p is similar to that of the asymptotically dS solutions previously considered [2]. In contrast, there exist solutions where w has no nodes, which are not seen in the asymptotically dS or Minkowski cases.

4 Dyon solutions

Dyon solutions to the EYM equations for a given negative Λ are found if we choose the adjustable shooting parameter a to be non-zero. Fig. 1 shows how the electric component, u, of the EYM equations starts at zero and monotonically increases to some finite value. The behavior of w, m , H, and p is similar to the monopole solutions.

Again we find a continuum of solutions for a continuous set of parameters a and b, where w crosses the axis an arbitrary number of times depending these parameters. Also similar to the monopole solutions is the existence of solutions where w does not cross the axis. This is in sharp contrast to the $\Lambda \geq 0$ case where the dyon solutions are forbidden.

5 Stability of the monopole solutions

The BK solutions and the dS-EYM solutions are unstable [7, 8, 9]. In contrast, the AdS black hole solutions [4] with $u = 0$ and the monopole solutions without nodes[5] are stable against spherically symmetric linear fluctuations.



Figure 1: Monopole and dyon solutions for $\Lambda = -0.01$ and $v = 1$. $(a, b) = (0, 0.001)$ and $(0, 0.005)$ for the monopole solutions and $(a, b) = (0.003, 0.001)$ and $(0.002, 0.0005)$ for the dyon solutions.

In order to derive the time dependent EYM equations, we use the most general expression for the spherically symmetric SU(2) gauge fields in the singular gauge:

$$A = \frac{1}{2e}\Big\{u\tau_3 dt + \nu\tau_3 dr + (w\tau_1 + \tilde{w}\tau_2)d\theta + (\cot\theta\tau_3 + w\tau_2 - \tilde{w}\tau_1)\sin\theta d\phi\Big\}, \tag{7}$$

where τ_i (i = 1,2,3) are the usual Pauli matrices and u, ν, w and $\tilde{w}$ depend on r and t. The boundary conditions $u = \nu = 0$ and $w^2 + \tilde{w}^2 = 1$ at $r = 0$ ensure regularity at the origin. Linearized equations when $u(t, r) = 0$, for the gauge fields $\delta w(t, r)$, $\delta\tilde{w}(t, r)$, $\delta\nu(t, r)$, $\delta p(t, r)$, and $\delta H(t, r)$ have been derived in the literature [3]. Fluctuations decouple in terms of $\delta w(t, r)$, $\delta p(t, r)$, and $\delta H(t, r)$ which form even-parity perturbations, and in terms of $\delta\tilde{w}(t, r)$ and $\delta\nu(t, r)$ which form odd-parity perturbations.

The equation for parity-odd perturbations in $\beta = r^2 p \delta\nu / w$, where $\beta(t,r) = e^{-i\omega t}\beta(r)$, is [5]

$$\left\{ -\frac{d^2}{d\rho^2} + U_\beta(\rho) \right\} \beta = \omega^2 \beta \quad , \quad U_\beta = \frac{H}{r^2 p^2}(1+w^2) + \frac{2}{w^2}\left(\frac{dw}{d\rho}\right)^2 \tag{8}$$

where $d\rho/dr = p/H$. Volkov et al. showed that for the BK solutions there appear exactly n negative eigenmodes ($\omega^2 < 0$) if w has n nodes [8]. Their argument applies to the asymptotically AdS case without modification. One concludes that the solutions with nodes in w are unstable against parity-odd perturbations.

For parity-even perturbations, where $\delta w(t,r) = e^{-i\omega t}\delta w(r)$, we find the equation

$$\left\{ -\frac{d^2}{d\rho^2} + U_w(\rho) \right\} \delta w = \omega^2 \delta w \quad , \quad U_w = \frac{H(3w^2-1)}{p^2 r^2} + 4v\frac{d}{d\rho}\left(\frac{Hw'^2}{pr}\right) \ . \tag{9}$$

Although the potential $U_w(\rho)$ is not positive definite, it is regular in the entire range $0 \leq \rho \leq \rho_{\max}$. The first term in U_w becomes negative for $w^2 < 1/3$. The Schrödinger equation (9) was solved numerically [5]. The potential for the solutions with $(a,b) = (0, 0.001)$, which has no node in w, has the lowest eigenvalue ω^2 of 0.028 and for $(a,b) = (0, 0.005)$, which has one node, has the lowest eigenvalue of 0.023. Therefore, these solutions are stable against parity-even perturbations, even if w has one node, and differ from the solutions where $\Lambda \geq 0$ where the parity-even perturbations are always unstable.

6 Conclusion

A continuum of new monopole and dyon solutions to the asymptotically AdS EYM equations have been found. In fig. 2 the spectrum of monopole solutions is plotted. The monopole solutions in which the magnetic component $w(r)$ of the gauge fields never vanishes, corresponding to the portion with $n = 0$ in fig. 2, were shown to be stable against linear perturbations. Solutions with $b \geq 0.7$ develop an event horizon, becoming black holes. The end point of the $n = 3$ portion of the curve in fig. 2 shows where the solutions become black holes.

Acknowledgments

This work was supported in part by the U.S. Department of Energy under contracts DE-FG02-94ER-40823 and DE-FG02-87ER40328.



Figure 2: The Q_M-$Mass$ plot of monopole solutions at $\Lambda = -0.01$. The number of nodes, n, in $w(r)$ is also indicated.

References

[1] R. Bartnik and J. McKinnon, Phys. Rev. Lett. **61** , 141 (1988).

[2] M.S. Volkov, N. Straumann, G. Lavrelashvili, M. Huesler and O. Brodbeck, Phys. Rev. D **54** 7243 (1996); T. Torii, K. Maeda and T. Tachizawa, Phys. Rev. D **52** R4272 (1995).

[3] M. S. Volkov and D. Gal'tsov, hep-th/9810070 (1998).

[4] E. Winstanley, Class. Quant. Grav. **16**, 1963 (1999).

[5] J. Bjoraker and Y. Hosotani, gr-qc/9906091 (1999).

[6] E. Witten, Phys. Rev. Lett. **38** 121 (1977).

[7] N. Straumann and Z. Zhou, Phys. Lett. **B237**, 353 (1990); Z. Zhou and N. Straumann, Nucl. Phys. **B360**, 180 (1991).

[8] M.S. Volkov, O. Brodbeck, G. Lavrelashvili, and N. Straumann Phys. Lett. **B349** 438 (1995).

[9] O. Brodbeck, M. Huesler, G. Lavrelashvili, N. Straumann and M.S. Volkov, Phys. Rev. D **54** 7338 (1996).